

call - 618
FAP-17

FAP 17

Fisheries Studies
and
Pilot Project (26)



FINAL REPORT

(Draft)

JUNE 1994



special
study



Supporting Volume
No. 24



INVESTIGATION OF PESTICIDE RESIDUE LEVELS IN FLOODPLAIN FISH IN BANGLADESH

ODA

Overseas Development Administration, U.K.

Special Study

FAP 17
FINAL REPORT

SUPPORTING VOLUME NO. 24

** Draft **



Investigation of Pesticide
Residue Levels in Floodplain
Fish in Bangladesh

A-4a

FAP 17
FISHERIES STUDIES
AND PILOT PROJECT

MAF-2111
22-12



June, 1994

Funded by ODA in conjunction with the Government of Bangladesh

ACKNOWLEDGEMENTS

The study was carried out jointly by FAP 17 and IFRB. The cooperation and effort of laboratory staff at IFRB are greatly appreciated and particular thanks go to Dr. N Chowdhury, Director and Dr. M Matin, Chief Scientific Officer. Thanks also go to Dr. A Hossain (Envirocare Ltd) who arranged visits to various scientific institutions and liaised with IFRB on behalf of FAP 17.

Close cooperation was provided by the Department of Fisheries and special thanks go to Mr R Banik, Senior Scientific Officer who collected data on pesticide use at each sampling site. FAP 20 kindly provided background data on several *beels* in the CPP from which our site selection was made. Finally, thanks go to Mr. C Price (BAFRU) who, despite a 9 hour flight delay managed to deliver frozen fish samples to the UK for further analysis at NRI, UK.

TABLE OF CONTENTS

| | Page No. |
|-------------------|----------|
| EXECUTIVE SUMMARY | iii |
| 1 INTRODUCTION | 1 |
| 1.1 Study Area | 2 |
| 2 METHODS | 4 |
| 2.1 Field | 4 |
| 2.2 Laboratory | 5 |
| 3 RESULTS | 6 |
| 4 DISCUSSION | 10 |
| 5 CONCLUSIONS | 12 |
| 6 RECOMMENDATIONS | 13 |
| REFERENCES | 14 |
| APPENDIX I | 16 |





LIST OF FIGURES

| Figure No. | Titles | Page No. |
|------------|----------------------------|----------|
| 1.1 | Location of sampling sites | 3 |

LIST OF TABLES

| Table No. | Titles | Page No. |
|-----------|--|----------|
| 1.1 | Pesticides used during 1992 to protect IRRI rice crop at selected sampling sites | 2 |
| 2.1 | Fish samples collected for pesticide residue analysis | 4 |
| 3.1 | Pesticide residue levels in fish (analysis by IFRB, AEC Savar, Dhaka) | 8 |
| 3.2 | Pesticide residue levels in fish (analysis by NRI, UK.) | 9 |
| 3.3 | Comparison of the results of pesticide residue analysis from IFRB (AEC), Bangladesh and NRI, UK. | 9 |

EXECUTIVE SUMMARY

In May 1992 samples of four fish species were collected from four small perennial *beels* in the Tangail Compartmentalisation Pilot Project area. The *beels* were surrounded by HYV rice fields receiving organophosphate and carbamate pesticides in March and April 1992. Pesticide residue analyses carried out in Bangladesh detected no organophosphate residues in fish tissue but revealed low levels of organochlorine pesticides: DDT, dieldrin and endosulphan. Further analyses undertaken in the UK as a cross-check of the results revealed low levels of DDD and DDE, metabolites of DDT, and also dieldrin but no endosulphan. The results indicated slight organochlorine pesticide pollution in floodplain ecosystems. The implications of the results relating to fish survival and human health are discussed.

1 INTRODUCTION

One of the principal aims of flood control and drainage schemes in Bangladesh is not only to provide protection for people and infrastructure against annual monsoon flooding but also to convert seasonal wetlands to agricultural land for increased rice production. In recent years reliance has been placed on high yield varieties (HYV) of rice which generally require greater amounts of pesticides and fertilisers to maintain production targets.

Pesticide use is not effectively controlled in Bangladesh and toxic persistent pesticides such as DDT, although banned from agricultural use, are reported to be imported illegally from India and also obtained from sources within country for protection of rice crops. The amount of registered pesticides used in Bangladesh has increased considerably in recent years. These comprise mainly organophosphates such as diazinon, dimecron and nogos used to protect HYV boro rice (Anon, 1989). There is growing concern about their possible harmful effects on floodplain fish and people (Abul Bashar & Reazuddin, 1990; World Bank, 1991). The situation is unlikely to improve in the absence of adequate monitoring programmes.

A literature search revealed only two previous studies on pesticide residue levels in floodplain ecosystems in Bangladesh, both of which were supported by the British Overseas Development Administration (Roberts *et al.*, 1986 and 1989). These studies were carried out in connection with surveys of ulcerative disease in fish, but included very few samples from floodplain habitats receiving pesticides. It was therefore decided that as part of the FAP 17 fisheries programme a preliminary investigation of pesticide residue levels in floodplain fish should be undertaken in areas known to receive a range of pesticides used to protect HYV rice in the winter season. However, since FAP 17 did not have the capability to carry out pesticide residue analyses, a survey was first made of laboratory facilities and technical expertise in various scientific institutions to assess the feasibility of undertaking such analyses in Bangladesh.

Details of institutions visited during an earlier FAP 17 Design Study (July-September, 1991) and as part of FAP 17 (January-April, 1992) are given in Appendix I. Of these only the Institute of Food and Radiation Biology (IFRB) of the Atomic Energy Research Establishment at Savar had both laboratory facilities and technical expertise to immediately undertake pesticide residue analysis. The Institute had previously carried out DDT analyses



in dried fish and had examined levels of a range of organochlorine and organophosphate pesticides in agricultural crops.


1.1 Study Area

The Tangail Compartmentalisation Pilot Project (CPP) was selected as the study area (Figure 1.1). The area surrounds the town of Tangail and lies within a flood control scheme designed to prevent annual floods from the Northern Dhaleswari and Pungli rivers. The area contains about 15 small perennial *beels* all of which are surrounded by HYV irrigated rice fields during the dry season. Based on information supplied by FAP 20, four *beels* were selected, ranging in size from 8 to 100 acres of water in the dry season. The areas immediately surrounding these *beels* planted with HYV rice ranged from 260 to 564 acres. Interviews with 10 farmers growing IRRI rice next to the *beels* revealed that a range of pesticides, mainly organophosphates, had been applied to protect the crop during March and April 1992 (Table 1.1). None of the farmers reported knowingly using DDT or any other organochlorine pesticide. No fishermen or farmers reported fish mortalities in the *beels* resulting from pesticide use in adjacent rice fields but fish from all *beels* suffered from ulcerative disease in 1991.

Table 1.1 Pesticides used during 1992 to protect IRRI rice crop at selected sampling sites

| Site | Pesticides Used | Pesticide Group | Months Applied |
|------------------------|--|-----------------------------------|----------------|
| Atia <i>beel</i> | Dimecron Nogos (= Dichlorvos) Furadan (= Carbofuran) | Organophosphate " Carbamate | March/April |
| Bhatchanda <i>beel</i> | Basudin (= Diazinon) | Organophosphate | April |
| Boro <i>beel</i> | Dimecron Diazinon Basudin (= Diazinon) | Organophosphate " " | March/April |
| Chakhar <i>beel</i> | Basudin Furadan (= Carbofuran) | Organophosphate Carbamate | March/April |

Figure 2.1 Location of sampling sites



2 METHODS

2.1 Field

Fresh fish samples were collected from catches of fishermen in four *beels* on 6 May 1992. The catches were wrapped in aluminium foil and stored in cool boxes containing ice made from deionised water. From the catches four species of fish were selected for pesticide residue analysis. The selected species included piscivores such as *taki* (*Channa punctatus*) and detritivores such the *baim* (*Mastacembelus armatus*) and *gutum* (*Lepidocephalus guntea*) both of which also prey on invertebrates in the mud. Details of the samples are given in Table 2.1. All fish were weighed, coded and labelled then stored in freezers in Tangail and Dhaka prior to dispatch to IFRB in Savar.

Table 2.1 Fish samples collected for pesticide residue analysis

| Site | Sample Code | Species | Numbers of Fish | Total Wt(g) | Length Range (cm) |
|-----------------|----------------|---|-----------------|-------------|-------------------|
| Atia beel | ODA-1 | <i>Mystus vittatus</i> (<i>tengra</i>) | 20 | 130 | 77-98 |
| Atia beel | ODA-3 | <i>Mastacembelus armatus</i> (<i>baim</i>) | 13 | 90 | 98-147 |
| Atia beel | ODA-5 | <i>Lepidocephalus guntea</i> (<i>gutum</i>) | 26 | 110 | 64-90 |
| Atia beel | ODA-4 + ODA-10 | <i>Channa punctatus</i> (<i>taki</i>) | 7 | 340 | 119-220 |
| Bhatchanda beel | ODA-15 | <i>Mastacembelus armatus</i> (<i>baim</i>) | 20 | 160 | 110-160 |
| Bhatchanda beel | ODA-16 | <i>Mystus vittatus</i> (<i>tengra</i>) | 20 | 160 | 80-101 |
| Bhatchanda beel | ODA-17 | <i>Channa punctatus</i> (<i>taki</i>) | 6 | 350 | 120-200 |
| Boro beel | ODA-7 | <i>Mastacembelus armatus</i> (<i>baim</i>) | 20 | 90 | 81-125 |
| Boro beel | ODA-8 + ODA-11 | <i>Lepidocephalus guntea</i> (<i>gutum</i>) | 20 | 80 | 66-92 |
| Chakhar beel | ODA-12 | <i>Lepidocephalus guntea</i> (<i>gutum</i>) | 20 | 110 | 70-96 |
| Chakhar beel | ODA-14 | <i>Mastacembelus armatus</i> (<i>baim</i>) | 20 | 140 | 105-132 |

2.2 Laboratory

Frozen fish samples were analysed at IFRB of the Atomic Energy Research Establishment in Savar near Dhaka. In all cases, samples were identified only by code numbers so as to avoid unintentional bias of the results. In all but one case, analyses were carried out on homogenised muscle tissue and not on organs such as the liver where residue levels would normally be expected to be higher. This approach was taken because fish of the size range analysed are usually gutted for human consumption and one of the aims of this investigation was to relate pesticide residue levels in fish to health risks for humans. Between 3 and 20 individual fish of each species, depending on the size range, were pooled to provide a single sample of 25 grammes for analysis. Thus, pesticide levels reported in this study represent averages for a number of individuals for each species.

Standard methods of extraction, clean-up and analysis using gas-liquid chromatography (GLC) were employed at IFRB to determine the concentration of a range of organochlorine and organophosphate pesticides. Various checks on the reliability of the methods were carried out at IFRB. These included "blind" replicate analyses on one sample (code ODA-16), duplicate analyses on other samples and the use of samples "spiked" with known concentrations of pesticide standards to determine recovery rates. As a further check on the reliability of these methods, three samples of muscle were sent to laboratories of the Natural Resources Institute (NRI), UK for analysis of organochlorines.



72

3 RESULTS

Results of analyses carried out at IFRB are presented in Table 3.1. A total of 10⁸ samples comprising 126 individual fish of four species was screened for organochlorine and organophosphate pesticide residues. Low concentrations of three types of organochlorine pesticides were detected by GLC analysis, these comprised DDT, dieldrin and endosulphan. These pesticides were reported not to have been used by local farmers during 1992. In contrast, the organophosphate pesticides used by the farmers in March and April 1992 were not detected by IFRB but these normally degrade very rapidly in the environment in contrast to DDT and dieldrin which are highly persistent and can accumulate in certain tissues, especially those with a high lipid content (Muirhead-Thomson, 1987). IFRB lacked the analytical equipment needed for the detection of carbamates. Therefore information on the effect of Furadan applications by farmers is lacking.

There was generally close agreement between the results of duplicate analyses of DDT and dieldrin carried out at IFRB. Even when "blind" replicates were used, agreement was good in all but one sample (Table 3.1).

The results of analyses undertaken at NRI, on 3 fish muscle samples are shown in Table 3.2. These results are compared with those of IFRB for the same samples (Table 3.3). The analyses by NRI revealed no DDT but detected very low concentrations of DDT breakdown products, DDD and DDE. This difference may be expected if further metabolism, especially anaerobic metabolism, occurred during storage or transit of samples to the UK. In fact the flight to the UK was delayed for 9 hours but NRI reported that the samples still arrived fully frozen.

The NRI analyses revealed no trace of endosulphan I and II in contrast to the IFRB results which indicated low levels in two of the three test samples. In addition, dieldrin was found in only one of the three samples analysed by NRI at a concentration within the range reported by IFRB.

In all but one case analyses were carried out on muscle. In this one case, analyses were carried out by mistake on whole ungutted fish (*L. guntea*) from two combined sites (sample: ODA-5+8+11). Since organochlorines such as DDT and dieldrin accumulate more readily

20

in fat deposits, higher concentrations are normally found in organs such as the liver and ovaries. Therefore, higher levels should be expected in whole fish samples. However, the results revealed the opposite: whole fish samples showed the lowest levels of DDT and the third lowest levels of dieldrin (Table 3.1).

Analysis of spatial variation in pesticide residue levels was difficult since only one species (*baim*) was found in the catches at all sites. In addition, concentrations were generally very low and in the case of DDT and dieldrin many were near the lower limits of detection. Therefore small variations in concentrations between sites were probably meaningless.

Pesticide residue levels recorded by IFRB varied between different species, highest levels were found in *taki*, a predatory species at the top of the aquatic food chain where persistent pesticides would normally be expected to accumulate. However, the results from NRI relating to DDD and DDE levels showed the opposite trend with highest values found in the *baim* and lowest in *taki*.

Table 3.1 Pesticide residue levels in fish (analysis by IFRB, AEC Savar, Dhaka)

| Species | Sample code number | No. of fish in one sample | DDT (mg/kg) | Dieldrin (mg/kg) | Endosulfan Sulfate II (mg/kg) | Endosulfan I (mg/kg) | Organophosphorus compounds (mg/kg) |
|--------------------------------|--------------------|---------------------------|----------------------------|----------------------------|-------------------------------|----------------------|------------------------------------|
| <i>Mastacembelus armatus</i> | ODA - 3 | 13 | 0.0024 0.0018 0.0021 | 0.0012 0.0026 0.0019 | ND | ND | ND |
| <i>Mastacembelus armatus</i> | ODA - 7 | 8 | 0.0039 0.0022 0.0031 | 0.0014 0.0016 0.0015 | ND | ND | ND |
| <i>Mastacembelus armatus</i> | ODA - 14 | 7 | 0.0040 0.0046 0.0043 | 0.0062 0.0060 0.0061 | ND | ND | ND |
| <i>Mastacembelus armatus</i> | ODA - 15 | 6 | 0.0027 0.0028 0.0028 | 0.0018 0.0020 0.0019 | ND | ND | ND |
| <i>Lepidocephalus guntea</i> * | ODA - (5+8+11) | 20 | 0.0017 0.0005 0.0011 | 0.0018 0.0005 0.0012 | 0.0121 | Trace | ND |
| <i>Lepidocephalus guntea</i> | ODA - 12 | 13 | 0.0013 0.0013 0.0013 | 0.0008 0.0008 0.0008 | 0.0103 | Trace | ND |
| <i>Channa punctatus</i> | ODA - 17 | 3 | 0.0249 | 0.0054 0.0064 0.0059 | 0.2275 | 0.0544 | ND |
| <i>Channa punctatus</i> | ODA - (4+10) | 7 | 0.0171 | 0.0074 0.0060 0.0067 | 0.1495 | 0.0544 | ND |
| <i>Mystus vittatus</i> | ODA - 1 | 20 | 0.0061 | 0.0032 | 0.0743 | Trace | ND |
| <i>Mystus vittatus</i> | ODA - 16 | 19 | 0.0015 | 0.0010 | 0.0214 | Trace | ND |
| <i>Mystus vittatus</i> ** | ODA-X=16 | 19 | 0.0012 | 0.0007 | 0.0170 | Trace | ND |
| <i>Mystus vittatus</i> ** | ODA-Y=16 | 19 | 0.0048 | 0.0006 | 0.0186 | Trace | ND |

Notes: ND = Not detectable (limits of detection not given), * = Analyses carried out by mistake on whole ungutted fish, ** = "Blind" homogenised replicates provided by FAP

Table 3.2 Pesticide residue levels in fish (analysis by NRI, UK.)

| Sample: ODA 10, Species: <i>Channa punctatus</i> | | | |
|---|--------|---------------------|--------|
| mg kg ⁻¹ | | mg kg ⁻¹ | |
| p'p-DDD : | 0.005 | Dieldrin : | 0.003 |
| p'p-DDE : | 0.004 | Endosulphan I : | <0.001 |
| o'p-DDT : | <0.003 | Endosulphan II : | <0.001 |
| p'p-DDT : | <0.004 | | |
| Sample: ODA 12, Species: <i>Lepidocephalus guntea</i> | | | |
| mg kg ⁻¹ | | mg kg ⁻¹ | |
| p'p-DDD : | 0.007 | Dieldrin : | <0.001 |
| p'p-DDE : | 0.007 | Endosulphan I : | <0.001 |
| o'p-DDT : | <0.004 | Endosulphan II : | <0.001 |
| p'p-DDT : | <0.005 | | |
| Sample: ODA 15, Species: <i>Mastacembalus armatus</i> | | | |
| mg kg ⁻¹ | | mg kg ⁻¹ | |
| p'p-DDD : | 0.012 | Dieldrin : | <0.001 |
| p'p-DDE : | 0.012 | Endosulphan I : | <0.001 |
| o'p-DDT : | <0.002 | Endosulphan II : | <0.001 |
| p'p-DDT : | <0.004 | | |

- Notes: 1. The results are reported above as mg active ingredient per kg of fish ("wet weight"). No corrections have been made to the results to take moisture or fat content into account.
2. Where no residues were detected, a result of less than (<) the limits of detection of the analytical technique used are reported. These lower limits of detection also take into account the individual sample weights, hence the difference in some of the figures.

Table 3.3 Comparison of the results of pesticide residue analysis from IFRB (AEC), Bangladesh and NRI, UK.

| Pesticide | Sample code no. | IFRB (mg/kg) | NRI (mg/kg) |
|----------------|-----------------|--------------|-------------|
| DDT | ODA - 15 | 0.0028 | ND(<0.007) |
| | ODA - 12 | 0.0013 | ND(<0.009) |
| | ODA - 10 | 0.0171 | ND(<0.006) |
| Dieldrin | ODA - 15 | 0.0019 | ND(<0.001) |
| | ODA - 12 | 0.0008 | ND(<0.001) |
| | ODA - 10 | 0.0067 | 0.003 |
| Endosulphan I | ODA - 15 | ND | ND(<0.001) |
| | ODA - 12 | Trace | ND(<0.001) |
| | ODA - 10 | 0.0544 | ND(<0.001) |
| Endosulphan II | ODA - 15 | ND | ND(<0.001) |
| | ODA - 12 | 0.0103 | ND(<0.001) |
| | ODA - 10 | 0.1495 | ND(<0.001) |

Note: Limits of detection of methods used at NRI are given in parentheses when pesticides could not be detected.


4 DISCUSSION

The results of this study indicate low level pollution of floodplain habitats by organochlorine pesticides DDT and dieldrin. Some doubt remains concerning endosulphan pollution since its presence in fish tissues was not confirmed by analytical tests made in the UK. This compound is the least persistent in the environment of the organochlorine pesticides. The source of the pollution has not been confirmed in this study but it seems likely to originate from the use of unregistered, illegal pesticides by farmers to protect their rice crops. The detection of DDT and its metabolites DDD and DDE together with dieldrin in fish muscle tissue gives little indication of the time of application of these compounds onto the floodplain environment since they are capable of not only persisting and bioaccumulating in various organisms within the food chain but can also persist in soil, silt and mud for long periods of time (Muirhead-Thomson, 1971, 1987; WHO, 1989).

With regard to DDT pollution and its derivatives DDD and DDE, concentrations of 0.025 mg/kg found in the most contaminated fish, *taki* (sample ODA-17, Table 3.1) and *baim* (sample ODA-15, Table 3.2) were well below safe values for tissue levels: 0-0.2 mg/kg (EPA, 1977). The same is true for dieldrin levels with safe concentrations of 0.3 mg/kg.

In terms of human health, levels of DDT were well below the Codex Alimentarius maximum residue limit for meat, 7 mg/kg (FAO/WHO, 1978) and the US Food and Drug Administration action level for DDT in fish fillets, 5 mg/kg. Even the most contaminated fish would have been safe to eat given the Codex Acceptable Daily Intake of DDT residue limit of 0.005 mg/kg body weight. Thus, a person weighing 50kg could safely consume 0.25 mg/kg of DDT per day which is equivalent to 10 kg of the most contaminated fish (0.025 mg/kg) per day. This is an enormous amount compared with the estimated daily per capita fish consumption of about 30g.

A literature search revealed only two previous studies of pesticide residues in freshwater fish in Bangladesh (Roberts *et al.*, 1986 and 1989). The studies were ancillary parts of broader studies of epizootic ulcerative fish disease. The first study found no detectable levels of pesticide residues in the two snakeheads analysed. The later study found no organophosphate or carbamate residues in fish tissues but detected low levels (0.01-0.03 mg/kg) of the



organochlorines, dieldrin, p'p-TDE and α HCH. The studies reported no direct link between pesticide pollution and the incidence of ulcerative fish disease in Bangladesh and Southeast Asia.

The absence of organophosphate pesticide residues in fish tissues found in this study and reported in previous studies carried out in Bangladesh does not imply that there is no pollution problem threatening fish survival in rice fields and adjacent water bodies. Although these compounds are reported to degrade rapidly (within days) in the environment they are highly toxic to fish, even at low concentrations. The immense damage caused by pesticides to fish cultured in the rice fields of Southeast Asia and beyond has been well established (e.g. Arce and Cagauan, 1988). However, in Bangladesh rice-fish culture is still in an infancy stage (CARE, 1992) and very little is known about the size and species composition of wild fish populations in boro rice fields. In a survey of 610 farmers throughout Bangladesh, 71% reported seeing fish mortalities when they applied pesticides to their boro rice crops (Anon., 1989). Unfortunately, the study did not report the magnitude of fish mortalities observed by farmers.

In Bangladesh, efforts are currently being made to promote the reduction in pesticide use in rice fields by the adoption of rice-fish culture associated with integrated pest management techniques (CARE, 1992). Such efforts should be encouraged since they offer large-scale potential not only for decreased environmental pollution but also increased cultured fish production in the winter season.

27

5 CONCLUSIONS

1. The results revealed the presence of low concentrations of organochlorine pesticide residues in fish tissues which were not considered harmful to human consumers.
2. From a total of 14 scientific institutions visited during this study to assess the feasibility of undertaking pesticide residue analysis in Bangladesh, only one was able to do so but needed assistance from FAP 17 to provide certain consumable items.
3. In Bangladesh, there is a serious lack of quantitative information on the effects of pesticides on inland floodplain fisheries.
4. There is a large potential to increase rice-fish culture in Bangladesh but the widespread use of pesticides to protect rice crops may pose a constraint on development.

gm

6 RECOMMENDATIONS

1. On the basis of the preliminary results obtained in this study, it is not considered appropriate for FAP 17 to continue more detailed and extensive investigations into pesticide residue levels in floodplain fish which would require considerable additional financial support to the project.
2. There is a need to strengthen the institutional capability to undertake pesticide residue analysis in Bangladesh. This would involve the provision of analytical equipment with the necessary maintenance services and training to improve technical expertise.
3. More information is needed on the effects of pesticides on wild fish populations in Bangladesh. The Department of Fisheries and the Department of Agricultural Extension should use their existing field personnel to carry out systematic, quantitative surveys of fish mortalities caused by pesticides.
4. In view of the great potential for increased rice-fish culture in Bangladesh, it is recommended that the development of small-scale integrated pest management schemes such as that established by CARE be included as an integral part of future rice-fish culture projects. Such projects must carefully consider the role of rapidly degradable pesticides in their development programmes.

REFERENCES

- Abul Bashar S.H.M. and Reazuddin Md, 1990
Towards sustainable development: issues of environmental pollution in Bangladesh.
Background paper prepared for Natural Conservation Strategy - Bangladesh. Submitted
September 1990.
- Anonymous 1989
A baseline survey of the pesticides used in Bangladesh with particular reference to their toxic
effects on Indian Major carps.
Unpub. M.Sc thesis, BAU, Mymensingh, August 1989.
- Arce R. G. & Cagauan A.G., 1988
Overview of pesticides use in rice-fish farming in Southeast Asia.
Paper presented at the Rice-fish farming Workshop, Ubon, Thailand, 21-25 March 1988.
- CARE, 1992
NOPEST Pilot Project Results. Rice-fish cultivation, 1992 Boro Season.
- EPA, 1977
Quality Criteria for Waters
Environmental Protection Agency, Washington, DC, 256 pp.
- FAO/WHO, 1978
Guide to Codex Maximum limits for pesticide residues.
First Issue. Codex Alimentarius Commission. CAC/PR 1-1978, 209 pp, FAO, Rome.
- Muirhead-Thompson R.C., 1971
Pesticides and Freshwater Fauna. Academic Press, London and New York.
- Muirhead-Thompson R.C., 1987.
Pesticide Impact on Stream Fauna with special reference to macroinvertebrates.
Cambridge University Press.

Roberts R.J., Macintosh D. J., Tonguthai K., Boongaratpalin S., Tayaputch N., Phillips M. J. & Millar S.D., 1986.

Field and Laboratory investigations into ulcerative fish diseases in the Indo-Pacific Region. FAO Project TCP/RAS/4508.

Roberts R. J., Wootten R., MacRae I., Millar S. D. and Struthers W., 1989
Ulcerative disease survey in Bangladesh.

Final Report to the Government of Bangladesh and the Overseas Development Administration, UK.

World Bank, 1991.

Bangladesh Fisheries Sector Review.

Report No 8830-8D, World Bank, March 1991.

WHO, 1989

Environmental health criteria for DDT and its derivatives - environmental aspects.
Report of International Programme on Chemical Safety.



25

Appendix I

26

Institutions and personnel visited during a survey of institutional capability for pesticide residue analysis in Bangladesh (July 1991 - March 1992).

1. Atomic Energy Research Establishment (AERE), Institute of Food & Radiation Biology (IFRB), Savar.
Contacts: Dr. N. Choudhury, Director
Dr. M. Matin, Chief Scientific Officer
2. Bangladesh Agricultural Research Institute (BARI), Gazipur.
Contacts: Dr. M K Chowdhury, Senior Scientific Officer
3. Bangladesh Agricultural University (BAU), Mymensingh
Contacts: Dr. Wahab, Associate Professor
4. Bangladesh Council of Scientific Industrial Research (BCSIR), Dhaka.
Contacts: Dr. A. Rahman, Director
5. Fisheries Research Institute (FRI), Mymensingh
Contacts: Professor M. Aminul Islam, Director
6. Fisheries Research Institute (FRI) Riverine Laboratory, Chandpur
Contacts: Mr. U. Haroon, Senior Scientific Officer
7. Institute of Postgraduate Studies in Agriculture (IPSA), Salna, Gazipur
Contacts: Dr. E. Ahsan, Director
8. International Centre of Diarrhoeal Disease Research, Bangladesh (ICDDR), Dhaka.
Contacts: Professor D. Habte, Director
9. Ministry of Agriculture, Department of Agricultural Extension, Plant Protection Wing, Dhaka.
Contacts: Mr. A K M Azad, Entomologist

10. Ministry of Environment and Forest, Department of Environment, Environment and Water Pollution Laboratory, Dhaka
Contacts: Dr M A H Pramanik, Director General
11. River Research Institute, Faridpur
Contacts: Mr A K M Shamsul Hoque, Director General
12. University of Dhaka, Dept of Biochemistry
Contacts: Professor M. Rahman, Chairman
Dr. A. Hossain, Assistant Professor
13. University of Dhaka, Institute of Nutrition and Food Science
Contacts: Dr. S. Keramat Ali, Director
14. University of Dhaka, Dept. of Zoology
Contacts: Professor Abu Tweb Abu Ahmed, Chairman

